

- If you are viewing this course as a recorded course after the live webinar, you can use the scroll bar at the bottom of the player window to pause and navigate the course.
- This handout is for reference only. Non-essential images have been removed for your convenience. Any links included in the handout are current at the time of the live webinar, but are subject to change and may not be current at a later date.

No part of the materials available through the continued.com site may be copied, photocopied, reproduced, translated or reduced to any electronic medium or machine-readable form, in whole or in part, without prior written consent of continued.com, LLC. Any other reproduction in any form without such written permission is prohibited. All materials contained on this site are protected by United States copyright law and may not be reproduced, distributed, transmitted, displayed, published or broadcast without the prior written permission of continued.com, LLC. Users must not access or use for any commercial purposes any part of the site or any services or materials available through the site.

Technical issues with the Recording?

- Clear browser cache using [these instructions](#)
- Switch to another browser
- Use a hardwired Internet connection
- Restart your computer/device

Still having issues?

- Call 866-782-9924 (M-F, 8 AM-8 PM ET)
- Email customerservice@OccupationalTherapy.com

CONTINUED

The Spaced Retrieval Technique: A How-To for OTs

Megan L. Malone, M.A. CCC-SLP

CONTINUED

Learning Outcomes

- As a result of this course, participants will be able to:
 - Describe the spaced retrieval technique and the evidence to support its use in rehabilitation contexts.
 - Provide two examples of goals that could be addressed using the spaced retrieval technique in occupational therapy.
 - Describe the process of how spaced retrieval is implemented in a therapy session.

CONTINUED

Disclosures

- **Presenter Disclosure:** Financial: Megan Malone received an honorarium for presenting this course. Non-financial: Megan is the co-author of the treatment manual, “Here’s How To Treat Dementia”, published by Plural Publishing in 2013 for which she received royalties.
- **Content Disclosure:** This learning event does not focus exclusively on any specific product or service.
- **Sponsor Disclosure:** This course is presented by OccupationalTherapy.com.

Acknowledgements

- Information shared in this presentation is provided through the work of various organizations & professionals:
 - Menorah Park Center for Senior Living/Myers Research Institute, Beachwood, OH
 - State of New York, Department of Aging
 - Hearthstone Alzheimer’s Care
 - Northern Speech Services
 - National Institute on Aging
 - Retirement Research Foundation

CONTINUED

Dementia Review

- Dementia is not a specific disease
- Dementia is a descriptive term for a collection of symptoms that can be caused by a number of disorders that affect the brain.
- Alzheimer's disease accounts for 60 to 80 percent of cases. Vascular dementia, which occurs after a stroke is the second most common dementia type.

Alzheimer's Association 2020

Q2

CONTINUED

Research Tells Us...

- Dementia is the loss of mental functions involving thinking, memory, reasoning, and language to such an extent that it interferes with a person's daily living.
- Dementia is a group of symptoms that can include:
 - Language disturbances (e.g., aphasia, dysphasia, anomia)
 - Challenging behaviors (e.g., repetitive questioning, wandering)
 - Difficulties with activities of daily living (e.g., dressing, personal grooming)
 - Personality disorders (e.g., disengagement, aggressive behaviors)

Alzheimer's Association, 2020

CONTINUED

CONTINUED

Memory

- Memory is dependent on organizing incoming information (attention) and highly developed encoding skills
- Memory is critical to our ability to acquire language, develop high level thinking, and effectively make decisions

CONTINUED

Memory Stages

- Encoding, Storage, and Retrieval are interactive processes
- The ability of one process affects the quality of another
 - Good encoding makes for good retrieval later on
- Deficit in one stage can lead to a deficit in another

CONTINUED

CONTINUED


Memory Definitions

- Working Memory, Short Term Memory:
 - Ability to use information as it's being processed (remembering phone number)
 - Primarily affected first with Alzheimer's and other dementias
- Long Term Memory:
 - Information from short term memory that is retained permanently
 - Declarative and Procedural Memory
 - Procedural memory is relatively-spared through the progression of dementia (FOUNDATION FOR SPACED RETRIEVAL TECHNIQUE)
 - Long Term Memory can be affected by dementia in both storing information and retrieving it.

Q1, Q5

CONTINUED

Learning & Memory in Dementia: Model of Memory



Squire, 1994

CONTINUED

CONTINUED

Mistaken Beliefs About Dementia

- Individuals with dementia cannot learn or remember information
- Best way to care for persons with dementia is to make them comfortable, accept their idiosyncrasies, and be patient with them.

Q3

CONTINUED

Circumvent the Deficits

- Persons with dementia do have weaknesses in the areas of learning and memory BUT a number of strengths exist as well.
 - Ability to learn procedures
 - Ability to read
- Research has shown that the learning of information and its retention depends heavily on how it is presented.
- KEY: Be aware of the weaknesses but FOCUS ON THE STRENGTHS!!!

CONTINUED

CONTINUED

Behavioral Interventions for Dementia

- Can be Direct or Indirect
- Direct
 - When an OT or other professional intervenes directly with individuals or group using an intervention
- Indirect
 - OT or other professional trains caregivers in an intervention, modifies the environment, or develops activities to maximize function
- Spaced Retrieval can be both Direct and Indirect

Mahendra & Hopper et. al, 2008

CONTINUED

The Spaced Retrieval Technique

CONTINUED

The Spaced Retrieval Technique

- Spaced Retrieval (SR)
 - Technique used to help persons with cognitive impairments recall important information over progressively longer intervals of time.
 - First used to address face-name learning in non-impaired individuals
 - Has been used successfully with patients with Alzheimer's Disease, Traumatic Brain Injury, Parkinson's Disease, and Dementia related to HIV (Bourgeois et. al, 2001; Camp et. al, 2008; Neundorfer et. al, 2004; Malone et. al, 2007)
 - Is an effective tool that therapists can use to help clients reach their goals in rehab therapy and is billable and reimbursable.
 - Takes advantage of the procedural memory system and is success-based.

Q4, Q5

The Spaced-Retrieval Technique

- The goal of SR:
 - To enable individuals to remember information for long periods (days, weeks, months, years) so that they can achieve long-term treatment goals.
- Therapists teach clients strategies that compensate for memory impairments, using procedural memory, including reading and repetitive priming.
- In addition, SR uses external aids to compensate for memory.

Brush & Camp, 1998

CONTINUED

The Spaced-Retrieval Technique

- Begin with a prompt question for the target behavior and teach the client to recall the correct answer.
- When retrieval is successful, the interval preceding the next recall test is increased.
- If a recall failure occurs, the participant is told the correct response and asked to repeat it.
 - Errorless learning: minimization of error responses during the presentation of target stimuli (Sohlberg & Turkstra, 2011).
- The following interval length returns to the last interval at which recall was successful.

Q5

CONTINUED

Treatment: Spaced Retrieval

- Goal: “Client will independently recall location of daily schedule to complete ADL’s, improve attendance at & participation in meals, and engage with peers 90% of trials.”
 - Question: “Where should you look to find your daily schedule?”
 - Answer: “Look at my walker”



CONTINUED

CONTINUED

Treatment: Spaced Retrieval

- Trial 1 (0 Seconds): Client Responds CORRECTLY
- Trial 2 (10 Seconds): Client Responds CORRECTLY
- Trial 3 (30 Seconds): Client Responds CORRECTLY
- Trial 4 (1 Minute): Client Responds INCORRECTLY
- Therapist provides client with correct response (“”), asks the client the prompt question again, allows the client to respond, and returns to the interval at which the client was last successful.
- Trial 5 (30 Seconds): Client Responds CORRECTLY
- Trial 5 (1 Minute): Client Responds CORRECTLY
- Client continues session; Therapist then probes through other therapy activities to continue training/practice of desired skill.

Q6

CONTINUED

SR Screening Measure

- Complete Screening Process
 - Quick and Easy
 - Tests clients' responses to correctly recall a target name over 3 different time intervals (immediately after presentation, 10 seconds later, and 15 to 20 seconds after that)
 - Client has 3 trials at each time interval to recall the target name correctly to pass the screen
 - CAN FOLD SR SCREEN INTO INITIAL CLIENT EVALUATION/ADMISSION INTERVIEW

Q7

CONTINUED

continued

A. Spaced-Retrieval Screen

(At each Trial, score as follows:
Correct response = + Incorrect response = - and
Record observations / what the participant says that's useful)

Start the Screening by saying:
"Today we are going to practice learning how to remember things better. We'll start by practicing remembering the name of this person." [Show photograph]

Step 1 Say:
"Her name (first and last) is Peggy Bailey.
What is her name?" Trial 1 ___ 2 ___ 3 ___
Observations: _____

Incorrect: Wait for about a minute, chatting with the participant, and then do a second trial of **Step 1**.
If incorrect on a second trial of **Step 1**, wait for about a minute, chatting with the participant, then do a third trial of **Step 1**.
If incorrect on a third trial of **Step 1**, then PROCEED DIRECTLY TO "EXIT LINE."

[If correct at **Step 1** on any trial, say:
"That's right. I am glad that you remembered."
THEN, Go to **Step 2**.

Step 2 (SHORT DELAY -- about 10 seconds) Say:
"Good. I will give you more chances to practice as I am working with you today. Let's try again. What is her name?"
Trial 1 ___ 2 ___ 3 ___
Observations: _____

Incorrect: Record response and go back to **Step 1** for a second or third trial at **Step 1**.
If incorrect on a third trial of **Step 2**, then PROCEED DIRECTLY TO "EXIT LINE."

[Correct: Say:
"That's right. I am glad that you remembered."
THEN Go to **Step 3**.

Step 3 (LONG DELAY -- about 15-20 seconds) Say:
"You are doing well remembering her name for a longer period of time, and that's the idea. I would like you to always remember her name. I will be practicing this with you during therapy by asking you often. So, what is her name?"
Trial 1 ___ 2 ___ 3 ___
Observations: _____

Incorrect: Say:
"Actually, her name is Peggy Bailey. What is her name?"
Go back to **Step 2**.
If incorrect on a third trial of **Step 3**, then PROCEED DIRECTLY TO "EXIT LINE."

[Correct: Say:
"That's right you are remembering for a longer period of time. You did a great job remembering her name. We are going to continue practicing this later today."
If you reach this point, continue with the study protocol.
EXIT LINE: "Thanks for trying so hard. Let's work on something else now."
If the client has not mastered **Step 1**, **Step 2** and **Step 3** successfully, try again on another day using a card with the name printed on it.
If the client fails the second screen, start therapy WITHOUT doing more screening for the study.
DO NOT ENROLL a person in the study who has not passed this Spaced-Retrieval screen.

continued

SR Goals: Prompt Question/Answer Examples

- Disorientation
 - "Where do you live?" (Answer: Name of Facility)
 - "What is your room number?" (Answer: Room #)
 - "What is your address?" (Answer: Client's address)
- Repetitive Questioning
 - Dependent upon question being asked

CONTINUED

SR Goals: Prompt Question/Answer Examples

- Energy Conservation:
 - Question: "What should you do before you begin a task?"
 - Answer: "Gather everything I need"
- IADL:
 - Question: "How do you remember which medicines you take?"
 - Answer: "Look at my list"
- Adaptive Equipment:
 - Question: "How should you reach for items safely?"
 - Answer: "Use my reacher."

CONTINUED

What Happens After the First SR Session?

- Therapist documents patient response to treatment and longest time interval attained
- The therapist begins (Trial 1 of the next session) by asking the client the prompt question and seeing if the client is immediately able to give the correct response.
 - This provides the client with an opportunity to demonstrate recall since the last treatment session, which may be 24 hours or more.

Q8

CONTINUED

CONTINUED

What Happens After the First SR Session?

- If the client can recall the correct answer to the prompt question (and the associated behavior, if applicable) then training on the question can cease for that session.
- If the client cannot recall the correct response, the clinician provides the correct answer, asks the client the prompt question again, allows the client to respond to demonstrate immediate recall and then training should resume, returning to the last time interval the client correctly recalled the response to in the previous session.

CONTINUED

What Happens After the First SR Session?

- Subsequent session example:
 - At start of any session following the initial training session on a prompt question/response, the clinician should allow the client to demonstrate recall of the information by asking the prompt question.
 - Client has left sided weakness.
 - Prompt ? : "How should you put on a shirt?;
 - Trial 1: "Use my dressing chart."
- Response Correct: Reinforce action or complete action & discontinue training for remainder of session (may choose to "spot check" retention of response throughout but formal timing of trials not necessary)
- Response Incorrect: Say, "Actually, you use your dressing chart. (provide correct response); "How should you put on a shirt? (ask prompt question again). Client responds "Use my dressing chart." "Good. Let's look at your chart.". Let's keep practicing-return to last successful time interval attained in previous session (e.g. 8 min), continue SR training based on client's responses.

CONTINUED

CONTINUED

When is an SR goal considered mastered?

- If a client is able to correctly respond to the prompt question and/or perform the targeted strategy at the beginning of 3 consecutive therapy sessions, the goal is considered mastered.
- It is important to make sure that the client is consistently performing the targeted strategy or response before discharging the goal.

CONTINUED

How much SR training does a client usually need?

- The amount of training required by a client will vary.
- The number of sessions is dependent upon:
 - Level of cognitive impairment of an individual client
 - Frequency of the sessions
 - Number of goals are being addressed using SR.

Clients enrolled in more frequent SR treatment sessions (i.e. 5 days week vs. 2 days/week) are likely to attain their goals more quickly.

Spaced Retrieval Goals

- Goal possibilities are endless
 - SR goals are NOT written any differently than other goals (SMART goals; ICF) (CMS, 2020)
 - https://www.cms.gov/Medicare/Billing/TherapyServices/Downloads/Mapping_Therapy_Goals_ICF.pdf
- FUNCTIONAL GOAL = SR GOAL

Q10

Spaced Retrieval Goals

- Measurement of goal attainment can be determined by area of focus & what allows for best measurement of progress.
 - by percentage or number of trials (“80% of time”; 4/5 trials)
 - “Patient will recall and demonstrate energy conservation techniques to decrease fatigue 80% of opportunities during a given activity.”
- OR
- Recalling or demonstrating target response for a set number of sessions (3 sessions recommended)
 - “Patient will recall and demonstrate strategy of properly using grab bar to transfer to shower at the initial trial of 3 consecutive therapy sessions using SR”

Q10

SR & Documentation

- SR is considered to be a MODALITY or APPROACH that therapists may use to help clients reach their goals.
- SR does not fit one particular diagnosis category
 - Use the ICD 10 Code that corresponds to the goal area you are addressing

SR Decision Making

- Questions to ask yourself when preparing to begin SR with a client:
 - What are the strengths of the client? What are the weaknesses (physical impairment, vision, etc.)?
 - What are the challenging behaviors being exhibited?
 - What prompt question & response will be used and is it and the answer meaningful for the client?
 - What other staff/family members will be involved in the training/carryover?

Q9

SR: An Interdisciplinary Process

- Caregiver/Family Input:
 - Consult with family/caregivers for possible goal ideas
= INCREASES BUY IN AND COOPERATION
 - Work on incorporating the patient and family's personal goals if possible.
 - Share prompts/responses with caregivers/staff after patient has demonstrated consistent success to increase generalization.

Case Study

- 75 year-old male; resident of assisted living facility. Diagnoses include Parkinson's disease and CVA 3-months prior. Referred by physician to receive occupational therapy services through home health upon discharge to home.
- Patient experiencing cognitive decline, is at risk for falls, and is experiencing decline in independence in ADL/IADLs.
- Patient Goal: "To remain in home and as independent as possible"

CONTINUED

Case Study

- Examples of goal areas:
 - Dressing
 - Showering
 - Bed mobility
 - Manage freezing episodes when completing mobility-related activities to reduce fall risk
 - Completion of home exercise program
- SR goal: “Patient will recall and demonstrate strategies to manage freezing episodes during movement to reduce fall risk at the initial trial of 3 consecutive sessions using the spaced retrieval technique.”

CONTINUED

Thank you!
meganmalone1025@gmail.com

CONTINUED

References

- Alzheimer's Association (2020). What is Dementia? Accessed at <https://www.alz.org/alzheimers-dementia/what-is-dementia>
- Bourgeois, M.S., Camp, C., Rose, M., White, B., Malone, M., Carr, J., & Rovine, M. (2003). A comparison of training strategies to enhance use of external aids by persons with dementia. *Journal of Communication Disorders*, 36, 361-378.
- Brush, J.A. & Camp, C.J. (1998). *A therapy technique for improving memory: Spaced retrieval*. Beachwood, OH: Menorah Park Center for the Aging.
- Camp, C. (1999). *Montessori-based Activities for persons with dementia*. Volume 1. Beachwood, OH: Menorah Park Center for the Aging.
- Camp, C., Schneider, N., Orsulic-Jeras, S., Mattern, J., McGowan, A., Antenucci, V., Malone, M., Gorzelle, G. (2006). *Montessori-based activities for persons with dementia*. Volume 2. Beachwood, OH: Menorah Park Center for Senior Living.

References

- Camp, C. J., & Malone, M. L. (2008) Mise en œuvre d'interventions de récupération espacée auprès de personnes atteintes de la maladie d'Alzheimer. Cahiers de la Fondation Médéric Alzheimer, number 3.
- CMS (2020). PT, OT, and SLP Services and the International Classification of Functioning, Disability, and Health (ICF). Accessed at https://www.cms.gov/Medicare/Billing/TherapyServices/Downloads/Mapping_Therapy_Goals_ICF.pdf
- Hopper, T., Mahendra, N., Kim, E., Azuma, T., Bayles, K.A., Cleary, S.J., et al. (2005). Evidence-based practice recommendations for working with individuals with dementia: Spaced-retrieval training. *Journal of Medical Speech-Language Pathology*, 13(4), 27-34.
- Loehr, J. & Malone, M. (2013). *Here's how to treat dementia*. San Diego: Plural Publishing.

References

- Malone, ML, Skrajner, MJ, Camp, CJ, Neundorfer M, Gorzelle, GJ: Research In Practice II: Spaced-Retrieval, A Memory Intervention. *Alzheimer's Care Quarterly*. (2007); 8(1): 65-74.
- Neundorfer, M.M., Camp, C.J., Lee, M. M., Malone, M. L., Carr, J. R., & Skrajner, M. J. (2004). Compensating for cognitive deficits in persons aged 50 and over with HIV/AIDS: A pilot study of a cognitive intervention. *Journal of HIV/AIDS & Social Services*, 3(1), 79-97.
- Sohlberg, M.M., & Turkstra, L.S.(2011). *Optimizing cognitive rehabilitation: Effective instructional methods*. New York: Guilford Press.
- Squire, LR. Declarative and nondeclarative memory: multiple brain systems supporting learning and memory. In: Schacter, DL, Tulving, E, eds. *Memory Systems*. Cambridge, MA: MIT Press; 1994: 203-232.